

## **RESOLUTION OIV/OENO 382B/2010**

### **UPDATE OF THE OIV COMPENDIUM OF METHODS OF ANALYSIS OF SPIRITUOUS BEVERAGES OF VITIVINICULTURAL ORIGIN –Overall determination of phenolic compounds in spirituous beverages of vitivinicultural origin without added caramel**

THE GENERAL ASSEMBLY

IN VIEW OF article 2 paragraph 2b iv of the agreement dated 3 April 2001 by which the international organisation of vine and wine was founded,

IN VIEW OF the actions of the 2009-2012 OIV strategic plan, in particular those aiming to reorganise the publications relating to vitivinicultural methods of analysis

CONSIDERING the work of the sub-commission of methods of analysis

IN VIEW OF the « Compendium of International Methods of Analysis of Spirituous Beverages of vitivinicultural origin » adopted in 2009

HAS DECIDED to introduce the method for the overall determination of phenolic compounds in spirituous beverages of vitivinicultural origin into the "Compendium of international methods of analysis of spirituous beverages of vitivinicultural origin"

### **OVERALL DETERMINATION OF PHENOLIC COMPOUNDS IN SPIRITUOUS BEVERAGES OF VITIVINICULTURAL ORIGIN WITHOUT ADDED CARAMEL**

Type IV method

#### **1. Definition**

Folin-Ciocalteu assay measures the total quantities of phenolic compounds originating from wood present in barrel-aged spirits that haven't received any added caramel.

This assay is not specific to phenolic compounds (cf. principle). Caramel also reacts to the Folin-Ciocalteu reagent. However, in the case of wood-aged spirituous beverages, the vast majority of results are related to the presence of phenolic compounds derived from the oak wood (VIDAL and Al, 1991).

Folin-Ciocalteu phenolic compound content corresponds to the response to the result described below. This result is expressed in mg of gallic acid/l by calibration.

## 2. Principle

All the phenolic compounds are oxidized by the Folin-Ciocalteu reagent. This reagent consists of a mixture of phosphotungstic acid and phosphomolybdic acid which is reduced, during the oxidation of the phenolic substances, into a mixture of blue molybdenum and tungsten oxides.

The blue colouring produced has a maximum absorption of about 750–760 nm. It is proportional to the quantity of oxidized phenolic compounds.

## 3. Apparatus

### 3.1. Standard laboratory apparatus, and in particular:

- Temperature-controlled bath (70° C),
- spectrophotometer.

## 4. Reagents

### 4.1. Folin-Ciocalteu reagent

This reagent is available for purchase ready to use. It can be prepared in the following manner:

- 100 g of sodium tungstate (No. CAS : 13472-45-2),
- 25 g of sodium molybdate (No. CAS : 7631-95-0), are dissolved in 700 ml of distilled water (No. CAS : 7732-18-5). Add:
- 50 ml of 85% phosphoric acid (No. CAS : 7664-38-2) ( $\rho_{20}=1.71$  g/mL),
- 100 ml of concentrated chlorhydric acid (No. CAS : 7647-01-0) ( $\rho_{20}=1.19$  g/mL).

Bring to a boil under reflux for 10 hours, then add:

- 150 G of lithium sulphate (No. CAS : 10377-48-7),
- a few bromine drops (No. CAS : 7726-95-6),

and bring to a boil for another 15 minutes. Cool and add 1 litre of distilled water.

#### **4.2. Anhydrous sodium carbonate.**

Prepare a 4.25% solution (m/v) in distilled water.

#### **4.3. Anhydrous gallic acid (No. CAS : 149-91-7),**

### **5. Procedure**

#### **5.1. Calibration in gallic acid**

Produce a hydroalcoholic gallic acid stock solution by weighed quantity, then some surrogate solutions by dilution (at least 2). The calibration range also includes a blank (hydroalcoholic solution). As an example, the range can include the following levels: 0.200 and 400 mg/L. Check the linearity of the calibration.

#### **5.2. Preparation of the samples**

The sample must be perfectly limpid and free of beeswing.

#### **5.3. Reaction**

In a test tube, introduce:

- 0.2 ml of sample (or of calibration solution)
- 1 ml of Folin-Ciocalteu reagent,
- 18.8 ml of sodium carbonate solution.

After stirring, bring to approximately 70°C for 20 minutes in the temperature-controlled bath, then cool under running cold water.

#### **5.4. Absorbance measurement at 760 Nm.**

Absorbance at 760 nm is measured under a 1 cm optical path.

### **6. Expression of results**

Express the result in mg of gallic acid/L (linear calibration), accounting for the possible dilution of the sample. If the absorbance is greater than 1, a new

measurement is carried out after dilution of the sample, if linearity is guaranteed.

## 7. References

1. BLOUIN J., LLORCA L., MONTREAU F.R., DUFOUR J.H., 1972. Etude des conditions optimales pour la détermination des composés phénoliques totaux par le réactif de Folin-Ciocalteu. *Connaissance de la vigne et du vin*, 6, 405-413.
2. VIDAL J.-P., CANTAGREL R., FAURE A., BOULESTEIX J.M., 1991. Comparaison de trois méthodes de dosage des composés phénoliques totaux dans les spiritueux. *FV OIV n°904*.